

Nearby Doorways, Parity Doublets and Parity Mixing in Compound Nuclear States

N. Auerbach,⁽¹⁾ J.D. Bowman⁽²⁾ and V. Spevak⁽¹⁾

⁽¹⁾*School of Physics and Astronomy, Tel Aviv University, Tel Aviv 69978, Israel*

⁽²⁾*Los Alamos National Laboratory, Los Alamos, New Mexico 87545*

(August 17, 1994)

Abstract

We discuss the implications of a doorway state model for parity mixing in compound nuclear states. We argue that in order to explain the tendency of parity violating asymmetries measured in ^{233}Th to have a common sign, doorways that contribute to parity mixing must be found in the same energy neighbourhood of the measured resonance. The mechanism of parity mixing in this case of nearby doorways is closely related to the intermediate structure observed in nuclear reactions in which compound states are excited. We note that in the region of interest (^{233}Th) nuclei exhibit octupole deformations which leads to the existence of nearby parity doublets. These parity doublets are then used as doorways in a model for parity mixing. The contribution of such mechanism is estimated in a simple model.

24.80.Dc, 24.60.Dr, 25.40.Ny, 25.85.Ec, 24.10.-i

Recent experiments on parity violation in compound nuclear (CN) states [1,2] are providing new information on the parity non-conserving (PNC) interaction. The experimental results for ^{232}Th showed that measured PNC asymmetries fluctuated about non-zero average, in contradiction to the purely random behavior expected on the basis of the statistical model of the CN. The understanding of PNC phenomena challenges theory to treat simultaneously the chaotic and the regular aspects of the CN system. A number of attempts were made to explain the new results in the framework of statistical models [3] or in models that combine nuclear dynamics with the statistical aspects [4,5]. Among the latter the doorway state approach was used in several theoretical works [4,5]. In Ref. [4] the spin-dipole (SD) resonance was used as the doorway to describe the PNC spreading width of the compound resonances. Later the same model was applied [6] in the calculation of the average longitudinal asymmetry. In Ref. [5] the $s_{1/2}$ and $p_{1/2}$ single-particle states were used as the doorways in an attempt to explain the constancy of signs. The common feature of these models is that they deal with distant doorways and involve only the one-body part of the PNC interaction. The term “distant doorways” refers to the fact that the position of the doorways is removed by $1\hbar\omega \sim 7$ MeV from the CN resonances under consideration. The model in Ref. [4] which involved the collective effects of the spin-dipole resonance was able to account for the PNC spreading width when a reasonable [7] value for the one-body PNC matrix element was used. However, when the same size matrix element was used the average asymmetry was almost two orders of magnitude smaller than measured. Recently more extensive theoretical investigations characterized several terms that could contribute to PNC asymmetries in the CN [8,9]. Additional terms were derived and evaluated. However these theories did not identify new terms that were large enough to explain the large non-fluctuating asymmetry.

The quantity measured in the experiments with polarized epithermal neutrons is the longitudinal asymmetry

$$P(E_r) = \frac{\sigma^+(E_r) - \sigma^-(E_r)}{\sigma^+(E_r) + \sigma^-(E_r)}, \quad (1)$$

where σ^+ , σ^- are the resonance part of the total cross sections for neutrons with positive

and negative helicities respectively. The index r designates the fact that the scattering is to a compound resonance $|r\rangle$ at energy E_r , carrying the quantum numbers $J = \frac{1}{2}^-$. We will refer to these resonances as $p_{1/2}$. The leading term of the asymmetry P can be written [6,9]

$$P(E_r) = -2 \sum_s \frac{\langle s | V^{\text{PNC}} | r \rangle \gamma_s}{E_r - E_s} \frac{\gamma_r}{\gamma_r}, \quad (2)$$

where γ_r and γ_s denote the escape amplitudes from resonances $|r\rangle$ and $|s\rangle$ due to the strong interaction force and are given by:

$$\begin{aligned} \gamma_r &= \langle \Phi_p^{(+)} | H | r \rangle, \\ \gamma_s &= \langle \Phi_s^{(-)} | H | s \rangle, \end{aligned} \quad (3)$$

where $\Phi_p^{(+)}$ and $\Phi_s^{(-)}$ denote the continuum p and s -wave functions in the elastic channels. The sum in Eq. (2) extends in principle over all states $|s\rangle$ that have the quantum numbers $J = \frac{1}{2}^+$. To reduce the sum in the equation one often introduces the doorway state approximation [10,11]. One seeks a subspace of states $|s\rangle$, denoted by $|d\rangle$ such that the coupling between these states and the states $|r\rangle - \langle r | V^{\text{PNC}} | d \rangle$ is sizeable or that the coupling between $|d\rangle$ and the continuum is strong or that both conditions are fulfilled so that when the sum in Eq. (2) is replaced by the partial sum over states $|d\rangle$ the result will be a good approximation to Eq. (2). In a formal treatment [9,10,12] one divides the space of states $|s\rangle$ into: $\{s\} = \{d\} + \{s'\}$. Assuming that the signs of the matrix elements $\langle r | V^{\text{PNC}} | s' \rangle$ are randomly distributed one derives [9–12] the following expression:

$$P(E_r) \simeq -2 \sum_d \frac{(E_r - E_d) \langle d | V^{\text{PNC}} | r \rangle \gamma_d}{[(E_r - E_d)^2 + \frac{\Gamma_d^2}{4}] \gamma_r}, \quad (4)$$

where γ_d is the escape amplitude of the doorway. Here Γ_d is the width of the doorway resulting from the coupling of $|d\rangle$ to the states $|s'\rangle$. The PNC matrix element is purely imaginary and the product of the two escape amplitudes is nearly pure imaginary because it involves opposite parities and thus the product: $(\langle d | V^{\text{PNC}} | r \rangle \gamma_r \gamma_d)$ is nearly real.

The PNC spreading width in the doorway approximation is given by [4]:

$$\Gamma_r^{\downarrow \text{PNC}} = \sum_d \frac{|\langle r | V^{\text{PNC}} | d \rangle|^2}{(E_d - E_r)^2 + \frac{\Gamma_d^2}{4}} \Gamma_d^{\downarrow}, \quad (5)$$

where $\Gamma_d^\downarrow$ is the spreading width of the doorway.

In the analysis of the recent class of experiments two basic quantities are determined [1,2]. Taking the ensemble of states $|r\rangle$ one determines the average asymmetry $\overline{P}$; the fluctuating part of the measured asymmetries yields the PNC spreading width in a given nucleus. In ^{233}Th the analysis [2] yields for $\Gamma_r^{\text{PNC}} \simeq 7.4 \times 10^{-7}$ eV and $\overline{P} \simeq (8 \pm 6)\%$. Equations (4) and (5) will be the basis of our discussion of the above two quantities.

Let us first apply these equations to the case of “distant” doorways. This was already the subject of study in Refs. [5] and [9] and we repeat this case in order to emphasize the difficulty one faces. Consider a one-body PNC potential of the form [13]

$$V^{\text{PNC}} = \epsilon 10^{-7} \frac{1}{2} \sum_i \{f(r_i), \sigma_i p_i c\} , \quad (6)$$

where $f(r)$ is some function of the distance r and the curly brackets denote the anticommutation operation. For this potential the doorway that will couple strongly to the state $|r\rangle$ is the giant spin-dipole resonance built on the state $|r\rangle$ i.e. by acting with the operator σr on $|r\rangle$. (For more details see Refs. [4,9]). The energy centroid of the spin-dipole resonance built on $|r\rangle$ is in ^{233}Th about 7 MeV above E_0 . The width of the spin-dipole is several MeV, and for the sake of an estimate let us take $\Gamma_{\text{SD}}^\downarrow = 3$ MeV. In Ref. [6,9] the ratio of the escape amplitudes in Eq. (4) was shown to be:

$$\frac{\gamma_{\text{SD}}}{\gamma_r} \simeq \frac{1}{\sqrt{N}} \frac{\sqrt{3}}{kR} \simeq \frac{1}{\sqrt{N}} \cdot 10^3 , \quad (7)$$

where $\frac{\sqrt{3}}{kR}$ is the ratio of penetrabilities and $N \sim 100$ is the number of particle-hole excitations that make up the collective spin-dipole in ^{232}Th . Using Eq. (4) and the value for $\overline{P}$ of 8% and the above values for the width and energy of the SD-doorway we find:

$$\langle r | V^{\text{PNC}} | \text{SD} \rangle = 1700 \text{ eV} . \quad (8)$$

If we use now Eq. (5) to estimate the collective PNC matrix element with the above values for E_d and $\Gamma_d^\downarrow$, we find that $\langle r | V^{\text{PNC}} | \text{SD} \rangle = 3$ eV. There is a discrepancy of almost three orders of magnitude with the value obtained in Eq. (8). The value of 3 eV is quite consistent with the predictions of the existing models for the PNC interaction in nuclei [7].

This leads us to the conclusion that one must go beyond the one-body aspects of the theory and the idea of distant doorways and address the question of the role of nearby doorways.

Let us consider the following physical scenario: the compound state which shows up in experiment as parity mixed, is close to a doorway so that the difference $|E_r - E_d| < \Gamma_d$ and that $\Gamma_d \simeq 100$ eV. Let us take for the sake of the estimate the value of $|E_r - E_d| = \Gamma_d/2$. Let us also assume that the ratio γ_d/γ_r is as before 10^3 due to the penetrability effect. Using again Eq. (4) we find for the PNC matrix element the value $\langle r|V^{\text{PNC}}|d\rangle = 4 \times 10^{-3}$ eV while when using Eq. (5) we estimate for this matrix element the value $\langle r|V^{\text{PNC}}|d\rangle = 6 \times 10^{-3}$ eV. The two numbers are not inconsistent as they were in the previous “distant” doorway case. Of course this consistency does not serve as a prove, it is just an indication that the discrepancy can be resolved when there are nearby doorways that couple to the compound state through a PNC interaction.

Let us now develop this idea further. In various reactions with low energy protons or neutrons and with good resolution one observes in the excitation functions structures that have a width Γ_d that is intermediate between the single-particle $\Gamma_{s.p.}$ and the compound width Γ_r , $\Gamma_{s.p.} \gg \Gamma_d \gg \Gamma_r$ [11,14]. These intermediate resonances are usually non-overlapping and their spacings D_d are such that $D_{s.p.} \gg D_d > D_r$. Very fine resolution experiments show that the intermediate resonances are composed of a number of compound resonances and that the intermediate resonances are enhancements in the excitation function of the compound nuclear states.

In heavy nuclei in the vicinity of $A = 240$ intermediate structure was seen and studied extensively. The most striking and most intriguing example is the intermediate structure observed in neutron induced fission (n, f). The cross section seen in these reactions shows groupings of compound states that are enhanced. The width of each such enhanced bump is about $\Gamma_d = 200$ eV and the bumps are spaced about $D_d = 700$ eV. A very elegant explanation of this behaviour of the (n, f) excitation function was given in terms of a double hump fission barrier [15]. The intermediate resonances observed in the reaction are due to

states in the second more shallow and lower fission barrier from which fission is easier and therefore enhanced.

Let us now use this information about the intermediate structure to consider some models for parity violation in these heavy nuclei. Let us first consider a model which connects to the observation of intermediate structure in the (n, f) reaction. The transition amplitude for this process can be expressed in the following form:

$$T_{fn} = \sum_{q^+} \frac{\langle f^+ | H | q^+ \rangle \langle q^+ | H | \Phi_n^{(+)} \rangle}{E - E_{q^+} + \frac{i}{2} \Gamma_{q^+}}, \quad (9)$$

where by $|f^+\rangle$ we denoted a state of positive parity describing the fissioning process, $|\Phi_n^{(+)}\rangle$ the incoming neutron continuum wave function states and $|q^+\rangle$ are positive parity compound states. The fact that the (n, f) process shows intermediate structure at some excitation energies, while the (n, n) excitation function does not, indicates that the mixing is enhanced for some states $|q^+\rangle$ that are in the vicinity of the fissioning states $|f^+\rangle$.

The two-body effective PNC interaction, derived using the meson-exchange models [7] and the two-body nuclear force have both similar ranges. Let us therefore assume that the matrix elements $\langle f^+ | H | q^+ \rangle$ and $\langle r | V^{\text{PNC}} | f^+ \rangle$ are related.

One can write

$$T = \sum_r \frac{\langle \Phi_s^{(-)} | H | f^+ \rangle \langle f^+ | V^{\text{PNC}} | r \rangle \langle r | H | \Phi_p^{(+)} \rangle}{(E - E_r + \frac{i}{2} \Gamma_r)(E - E_{f^+} + \frac{i}{2} \Gamma_{f^+})}, \quad (10)$$

which is the PNC T matrix for neutron scattering in the doorway state approximation with the doorway being the fission state $|f^+\rangle$. The corresponding asymmetry is then

$$P(E_r) = -2 \frac{(E_r - E_{f^+}) \langle \Phi_s^{(-)} | H | f^+ \rangle \langle f^+ | V^{\text{PNC}} | r \rangle}{\langle \Phi_p^{(+)} | H | r \rangle [(E_r - E_{f^+})^2 + \frac{1}{4} \Gamma_{f^+}^2]}. \quad (11)$$

The enhancements observed in the (n, f) reaction will also occur in the asymmetry given in Eq. (11) if the compound state $|r\rangle$ is in the vicinity of the doorway $|f^+\rangle$.

It is remarkable that in ^{233}Th (as well as in ^{231}Th) one finds [15–17] that the energy potential as a function of deformation is triple humped. It has been shown [16] that the existence of the third minimum in the potential is related to the appearance of *pear-like*

octupole deformations in the body frame of reference. This spontaneous breaking of reflection symmetry implies the existence of nearby (a few tens of keV apart) parity doublets [18,19]. In fact at somewhat higher energies than considered here such doublets were observed in ^{233}Th in the (n, f) experiments [17]. It is worth mentioning that the states in the third minima with excitation energies $E_{\text{exc}} \sim 4 - 8$ MeV belong to the class of hyperdeformed states, since the quadrupole and octupole deformations are very large, of the order $\beta_2 \simeq 0.90$, $\beta_3 \simeq 0.35$ [20,16].

Consider now such a $J = \frac{1}{2}^\pm$ doublet denoted by $|f^\pm\rangle$. The energy difference $|E_{f^+} - E_{f^-}|$ is equal to a few tens of keV, which is of course small but larger than $|E_r - E_{f^+}|$ or Γ_{f^+} . Let us now single out the $|f^-\rangle$ component in the wave function of $|r\rangle$

$$|r\rangle = a_{rf^-}|f^-\rangle + |r'\rangle . \quad (12)$$

Then

$$\langle f^+ | V^{\text{PNC}} | r \rangle = a_{rf^-} \langle f^+ | V^{\text{PNC}} | f^- \rangle . \quad (13)$$

We drop the contribution of $|r'\rangle$ to the matrix element in the above equation because: (i) in view of the (n, f) experimental results the coupling of $|r\rangle$ to $|f^-\rangle$ should dominate, meaning that a_{rf^-} is relatively large and (ii) because of the close relation between $|f^+\rangle$ and $|f^-\rangle$ the PNC matrix element should also be relatively large. Substituting Eq. (13) into Eq. (11) we find:

$$\begin{aligned} P(E_r) &= -2 \frac{\langle f^+ | V^{\text{PNC}} | f^- \rangle (E_r - E_{f^+}) \langle \Phi_s^{(-)} | H | f^+ \rangle a_{rf^-}}{[(E_r - E_{f^+})^2 + \frac{1}{4}\Gamma_{f^+}^2] \langle \Phi_p^{(+)} | H | r \rangle} . \end{aligned} \quad (14)$$

In order to evaluate the average part of $P(E_r)$ let us proceed with this model and use Eq. (12) to estimate $\langle \Phi_p^{(+)} | H | r \rangle$. Recognizing as before that the component of $|f^-\rangle$ in $|r\rangle$ is relatively large and that $|f^-\rangle$ connects to exit channel as the (n, f) experiments show we may write

$$\langle \Phi_p^{(+)} | H | r \rangle \simeq a_{rf-} \langle \Phi_p^{(+)} | H | f^- \rangle . \quad (15)$$

Substituting this into Eq. (14) we find:

$$P(E_r) = -2 \frac{\langle f^+ | V^{\text{PNC}} | f^- \rangle (E_r - E_{f+}) \langle \Phi_s^{(-)} | H | f^+ \rangle}{[(E_r - E_{f+})^2 + \frac{1}{4} \Gamma_{f+}^2] \langle \Phi_p^{(+)} | H | f^- \rangle} . \quad (16)$$

We see that except for the energy difference in the numerator the sign of P is independent of $|r\rangle$ and is fixed. If the compound states $|r\rangle$ observed in experiment are located on one side of E_{f+} the asymmetries P will have the same sign. A change in sign should occur when the energies E_r cross E_{f+} . This is similar to the circumstances that occur in isobaric analog resonances [11,14].

Realizing the fact that we deal with parity doublets having similar structure we take the ratio of the two escape amplitudes from $|f^+\rangle$ and $|f^-\rangle$ to depend only on the penetrability factor which is about 10^3 . Taking from the experiment [2] the value $\overline{P} \simeq 8 \times 10^{-2}$ and using $|E_r - E_{f+}| = \Gamma_{f+}/2$ we derive from Eq. (16):

$$\frac{\langle f^+ | V^{\text{PNC}} | f^- \rangle}{\Gamma_{f+}} \simeq 4 \times 10^{-5} . \quad (17)$$

For $\Gamma_{f+} \simeq 200$ eV the PNC matrix element is $|\langle f^+ | V^{\text{PNC}} | f^- \rangle| \simeq 10^{-2}$ eV. This is not an unreasonable value for such matrix element. Recall that in Ref. [4,6] we estimate that single-particle configurations give PNC matrix elements of about 0.3 eV.

We should mention that the breaking of reflection symmetry in the body frame of reference in actinides $A \leq 229$ is well established. These nuclei exhibit [21] (see reviews [19]) at low-energies octupole deformations $\beta_3 \simeq 0 - 0.12$, enhanced E1 transitions and large dipole moments in the body frame of reference.

We also note that effects of parity violation were observed in fission reactions induced by polarized neutrons [22–24]. A theoretical discussion of this phenomenon was presented some years ago [23] in which the notion of octupole deformations and parity doublets was invoked. The connection between PNC effects in the (n, f) reactions and the PNC experiments discussed in the present work should therefore not be overlooked.

We wish to thank G.F. Bertsch, V.V. Flambaum, G.T. Garvey, W.C. Haxton, M.B. Johnson, A.K. Kerman and D.H. Wilkinson for helpful discussions. This work was supported in part by the US-Israel Binational Science Foundation. Part of this work was performed at the Institute for Nuclear Theory in Seattle.

REFERENCES

- [1] J.D. Bowman *et al.*, Phys. Rev. Lett. **65**, 1192 (1990); X. Zhu *et al.*, Phys. Rev. C **46**, 768 (1992).
- [2] C.M. Frankle *et al.*, Phys. Rev. Lett. **67**, 564 (1991); C.M. Frankle *et al.*, Phys. Rev. C **46**, 778 (1992).
- [3] A. Müller and H.L. Harney, Phys. Rev. C **45**, 1955 (1992); J.B. French, V.K.B. Kota, A. Pandey, and S. Tomsovic, Ann. Phys. (N.Y.) **181**, 198 (1988); O. Bohigas and H.A. Weidenmüller, Annu. Rev. Nucl. Part. Sci. **38**, 421 (1988); J.B. French and V.K.B. Kota, Annu. Rev. Nucl. Part. Sci. **32**, 35 (1982).
- [4] N. Auerbach, Phys. Rev. C **45**, R514 (1992).
- [5] J.D. Bowman *et al.*, Phys. Rev. Lett. **68**, 780 (1992).
- [6] N. Auerbach and J.D. Bowman, Phys. Rev. C **46**, 2582 (1992).
- [7] B. Desplanques, J.F. Donoghue, and B.R. Holstein, Ann. Phys. (N.Y.) **124**, 449 (1980).
- [8] C.H. Lewenkopf and H.A. Weidenmüller, Phys. Rev. C **46**, 2601 (1992).
- [9] N. Auerbach and V. Spevak, Phys. Rev. C, in print.
- [10] N. Auerbach, J.Hufner, A.K. Kerman, and C.M. Shakin, Rev. Mod. Phys. **44**, 48 (1972).
- [11] H. Feshbach, A.K. Kerman, and R.H. Lemmer, Ann. Phys. (N.Y.) **41**, 230 (1967).
- [12] A.K. Kerman and A.F.R. de Toledo Piza, Ann. Phys. (N.Y.) **48**, 173 (1968).
- [13] F.C. Michel, Phys. Rev. **133**, B329 (1964).
- [14] A.K. Mekjian, Adv. Nucl. Phys. **7**, 1 (1973).
- [15] S. Bjørnholm and J.E. Lynn, Rev. Mod. Phys. **52**, 725 (1980).
- [16] V.V. Pashkevich, Nucl. Phys. **A169**, 275 (1971); P. Möller and J.R. Nix, in *Proc. Int.*

- Symp. on Physics and Chemistry of Fission*, Rochester, 1973 (IAEA, Vienna, 1974) Vol. 1, p.103; R. Bengtsson *et al.*, Nucl. Phys. **A473**, 77 (1987).
- [17] J. Blons *et al.*, Nucl. Phys. **A414**, 1 (1984); Nucl. Phys. **A502**, 121c (1989).
- [18] A. Bohr and B. Mottelson *Nuclear Structure*, (Benjamin, New York, 1975), Vol. 2.
- [19] I. Ahmad and P.A. Butler, Annu. Rev. Nucl. Part. Sci. **43**, 71 (1993); S. Åberg, H. Flockard and W. Nazarewicz, Annu. Rev. Nucl. Part. Sci. **40**, 439 (1990); W. Nazarewicz, Nucl. Phys. **A520**, 333c (1990).
- [20] S. Ćwiok *et al.*, Phys. Lett. **322B**, 304 (1994) and referenced therein.
- [21] G.A. Leander, W. Nazarewicz, G.F. Bertsch, J. Dudek, Nucl. Phys. **A453**, 58 (1986).
- [22] G.A. Petrov *et al.*, Nucl. Phys. **A502**, 297c (1989); A.Ya. Alexandrovich *et al.*, Nucl. Phys. **A567**, 541 (1994); F. Gönnenwein *et al.*, Nucl. Phys. **A567**, 303 (1994).
- [23] O.P. Sushkov and V.V. Flambaum, Sov. J. Nucl. Phys. **31**, 28 (1980); Phys. Lett. **94B**, 277 (1980).
- [24] V.E. Bunakov and V.P. Gudkov, Nucl. Phys. **A401**, 93 (1983).